


Article

Synthesis of β -Difluoroalkyl Azides via Elusive 1,2-Azide Migration

An unprecedented 1,2-azide migration has been developed through *gem*-difluorination of the easily available α -vinyl azides with *in situ*-generated $\text{PhIF}_2 \cdot \text{HF}$. This approach provides a platform for the downstream synthesis of β -difluorinated alkyl azides in high yields with broad substrate scope and excellent functional group tolerance. The DFT calculations suggest 1,2-azide migration occur via a three-membered azacyclic transition state.

Yongquan Ning, Paramasivam Sivaguru, Giuseppe Zanon, Edward A. Anderson, Xihe Bi

bixh507@nenu.edu.cn

HIGHLIGHTS

Rapid synthesis of β -difluorinated alkyl azides

Elusive 1,2-azide migration of α -vinyl azides

Late-stage modifications of natural products and drugs

Novel mechanism via a three-membered transition state

Article

Synthesis of β -Difluoroalkyl Azides via Elusive 1,2-Azide Migration

Yongquan Ning,¹ Paramasivam Sivaguru,¹ Giuseppe Zanoni,² Edward A. Anderson,³ and Xihe Bi^{1,4,5,*}

SUMMARY

The development of azide migration reactions is a formidable challenge because of the potential competition of side processes driven by the release of molecular nitrogen. Here, we report a conceptually novel 1,2-azide migration in an unprecedented *gem*-difluorination of the readily available α -vinyl azides, a transformation that enables the synthesis of a range of novel β -difluorinated alkyl azides. The practicality of the method is demonstrated by broad substrate scope, excellent functional group compatibility, and high yields. The migrating group selectivity can be tuned through electronic effects, and DFT calculations suggest 1,2-azide migration occurs via a three-membered azacyclic transition state. By using routine protocols, the β -difluorinated alkyl azide products can be easily transformed to biologically relevant β -difluorinated amines—common structural motifs in pharmaceuticals, thus demonstrating the utility of these fluorinated organic azides for pharmaceutical synthesis as well as other synthetically useful derivatives.

INTRODUCTION

The construction of carbon-nitrogen (C–N) bonds is one of the most important transformations in organic chemistry because of the prevalence of nitrogen-containing compounds in pharmaceuticals, natural products, and organic materials.^{1–3} Organic azides are widely recognized as convenient tools in such processes because of their superior ability to participate in a variety of C–N bond forming reactions.^{4–12} Among their many applications,^{13,14} organic azides are easily transformed into amines, and readily undergo C–N bond formation via nitrogen-centered reactive intermediates (e.g., nitrenoid chemistry, driven by the release of molecular nitrogen),^{15–18} or as a 1,3-dipole (e.g., click reactions).^{12,15} In contrast, transformations in which the azide functionality remains intact in the final product are almost unknown, the sole exception being a pericyclic 1,3-allylic azide rearrangement,^{19–24} which is limited in utility because of the generation of mixtures of allylic azide products (Figure 1A). The development of new azide migration reactions would therefore be of great value, not only because they would provide an ideal approach to organic azides inaccessible by existing methods but also due to the conceptual challenge of avoiding the highly entropically favorable loss of N₂.^{13,14}

β -difluoroalkylamines are key structural motifs in a variety of medically important compounds such as anticancer, anticholinergic, and anti-inflammatory therapeutic agents (Figure 1B),^{25–31} however, current synthetic approaches require tedious multi-step operations.²⁵ We questioned whether the readily available α -vinyl azides—a class of structurally unique functionalized alkene that displays a rich reactivity profile⁶—could offer a much more rapid entry to β -difluoroalkylamines

The Bigger Picture

The development of azide migration reactions is a formidable challenge, and the only sole exception is the pericyclic 1,3-allylic azide rearrangement. This is probably because of the liable decomposition of azides by the release of molecular nitrogen. In this article, we introduce a conceptually novel 1,2-azide migration through *gem*-difluorination of the readily available α -vinyl azides mediated by Py·HF and PIDA, allowing for the synthesis of β -difluorinated alkyl azides. This novel rearrangement reaction displays high reaction efficiency, broad substrate scope, and excellent functional group tolerance. Further transformations of these fluorinated azide products could afford a variety of *gem*-difluorinated nitrogen-containing molecules of potential interest in medicinal chemistry. Therefore, the discovery described here represents a conceptual advance in azide preparation and has opened up an avenue to the azide migration highly desired by synthetic chemists.






Figure 1. Development of 1,2-Azide Migration Reactions

(A) Applications of organic azides in organic synthesis.

(B) Examples of bioactive molecules containing a *gem*-difluoroalkylamine motif.

(C) Planned *gem*-difluorination of vinyl azides with *in situ*-generated $\text{PhIF}_2\cdot\text{HF}$ via an unprecedented 1,2-azide migration process (this work).

via a *gem*-difluorination and 1,2-azide migration (Figure 1C). The related *gem*-difluorination and 1,2-aryl migration of styrenes, recently been described by Jacobsen^{32,33} and others,^{34–38} provides an efficient and powerful means to access the *gem*-difluoroalkyl motif, a useful bioisostere of common polar functional groups.³⁹ However, with the exception of isolated examples involving benzyl ether (OBn) and sulfonamide (NHTs) groups,⁴⁰ these rearrangements proceed almost exclusively via arene migration. Under similar oxidative fluorination conditions (pyridine·HF and hypervalent iodine),³⁶ we report herein a conceptually novel 1,2-azide migration, which enable the efficient synthesis of a wide range of *gem*-difluorinated alkyl azides (Figure 1C). The migratory preference can be tuned through electronic effects, enabling a product switch in the reactions of α -aryl vinyl azides (i.e., azide migration versus arene migration). Further, the product *gem*-difluoroalkylazides can be readily converted to the corresponding biologically relevant *gem*-difluoroalkylamines.^{41,42}

RESULTS AND DISCUSSION

We first sought to establish conditions for the *gem*-difluorination and 1,2-azide migration reaction by using vinyl azide **1a** as a model substrate (Table 1). Subjection of **1a** to oxidative fluorination conditions employed for *gem*-difluorination and 1,2-aryl migration of styrenes³⁶ (pyridine·HF as fluorine source, bis(acetoxy)iodobenzene (PIDA) as oxidant in CH_2Cl_2 at 25°C) afforded the 1,2-azide migration product **2a** in excellent yield (entry 1). An oxidant was essential (entry 2), whereas other potential fluorine sources (e.g., AgF ,⁴³ CsF ,⁴⁴ $\text{Et}_3\text{N}\cdot\text{HF}$ ⁴⁵) led to no reaction (entries 3–5). Changing the oxidant to the more potent bis(trifluoroacetoxy)iodobenzene (PIFA)³⁴ resulted in a significantly reduced yield of **2a** (30%, entry 6), whereas use of PhIO ⁴⁴ offered no advantage (entry 7). A brief survey of other solvents revealed poor or no conversion in acetonitrile and DMSO (entries 8 and 9); similarly, lowering the reaction temperature to -40°C or -78°C led to reduced yields (entries 10 and 11).

Having identified optimal conditions (Table 1, entry 1), the scope of this *gem*-difluorination/1,2-azide migration was first explored with a range of α -alkyl vinyl azides (Figure 2). As shown in Figure 2A, vinyl azides with primary alkyl (**1ab**, **1ac**, and **1ah**), secondary alkyl (**1ad**, **1ae**, **1af**, and **1ai**), or benzyl (**1ag**) groups were

¹Department of Chemistry, Northeast Normal University, Changchun 130024, China

²Department of Chemistry, University of Pavia, Viale Taramelli 12, Pavia 27100, Italy

³Chemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford OX1 3TA, UK


⁴State Key Laboratory of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, China

⁵Lead Contact

*Correspondence: bixh507@nenu.edu.cn

<https://doi.org/10.1016/j.chempr.2019.12.004>

Table 1. Optimization of the Reaction Conditions



| Entry | Oxidant | Fluorinating Agent | Solvent | T (°C) | Yield (%) ^a |
|-------|---------|----------------------|---------------------------------|--------|------------------------|
| 1 | PIDA | Py·HF | CH ₂ Cl ₂ | 25 | 92 |
| 2 | none | Py·HF | CH ₂ Cl ₂ | 25 | 0 |
| 3 | PIDA | AgF | CH ₂ Cl ₂ | 25 | 0 |
| 4 | PIDA | CsF | CH ₂ Cl ₂ | 25 | 0 |
| 5 | PIDA | Et ₃ N·HF | CH ₂ Cl ₂ | 25 | 0 |
| 6 | PIFA | Py·HF | CH ₂ Cl ₂ | 25 | 30 |
| 7 | PhIO | Py·HF | CH ₂ Cl ₂ | 25 | 90 |
| 8 | PIDA | Py·HF | CH ₃ CN | 25 | 47 |
| 9 | PIDA | Py·HF | DMSO | 25 | 0 |
| 10 | PIDA | Py·HF | CH ₂ Cl ₂ | -40 | 60 |
| 11 | PIDA | Py·HF | CH ₂ Cl ₂ | -78 | 30 |


Reaction conditions: **1a** (0.5 mmol), PIDA (0.6 mmol), and Py·HF (2.5 mmol) in CH₂Cl₂ (5 mL) at 25°C for 1 min.

^aIsolated yield.


smoothly converted to products **2ab–2ai** in 75%–95% yield. α -alkyl vinyl azides featuring a variety of functionalities on the terminal carbon of the alkyl chain (e.g., chloro, nitro, cyano, sulfonyl, ester, carboxylic acid, and ether groups) also proved to be suitable substrates, giving products **2aj–2au** in 62%–95% yield. This included tolerance of other potentially reactive azides on the alkyl chains (substrates **1ak** and **1au**), which afforded the rearrangement products **2ak** and **2au** in 80% and 85% yields, respectively. Substrates with α -substituents such as acrylic ester (**1ba**), cyclohexenyl (**1bb**), and internal alkynyl (**1bc**) moieties (Figure 2B) also selectively delivered the corresponding *gem*-difluorination and 1,2-azide migration products (**2ba–2bc**) in good yield, without detrimental fluorination of the other unsaturated functionality of the α -substituents. These intriguing results clearly reveal that if the compound contains two unsaturated groups, olefins bearing an azide group have higher reactivity than free unsaturated groups.

We applied this methodology to the late-stage modification of natural product derivatives bearing a vinyl azide (Figure 2C). Substrates derived from menthol (**1ca**), vitamin E (**1cb**), and diacetone-D-galactose (**1cc**) afforded the desired products **2ca–2cc** in 62%–89% yields. These examples highlight the potential utility of this method in the straightforward derivatization of bioactive molecules. This difluoroazidation methodology can also be used to build polyfunctionalized products; as depicted in Figure 2D, 2,7-diazidoocta-1,7-diene (**1da**) and 2,8-diazidonona-1,8-diene (**1db**) provided the bis-difluoroazides, **2da** and **2db**, in 84% and 85% yield, respectively. Even a substrate containing four vinyl azide units (**1dc**) was converted to the desired product (**2dc**) in good yield (75%). Such azides are potentially useful building blocks for the synthesis of linear and poly(triazole) dendrimers via click chemistry, opening up applications in material science and medicinal chemistry.^{46–48}


Spurred on by the positive results obtained with α -alkyl vinyl azides, and with previous reports on styrene *gem*-difluorination/1,2-aryl migration in mind,^{32–38} we next




A 1,2-Azide migratory *gem*-difluorination of alkyl-substituted vinyl azides



B Selective 1,2-azide migratory *gem*-difluorination of vinyl azides bearing an unsaturated group



C 1,2-Azide migratory *gem*-difluorination of natural product derivatives



D 1,2-Azide migratory *gem*-fluorination of dimeric and tetrameric vinyl azides




Figure 2. Scope of 1,2-Azide Migratory *gem*-difluorinations

- (A) Varying the α -alkyl group.
(B) Unsaturated substrates.
(C) Natural product derivatives bearing a vinyl azide moiety.
(D) Polyazide substrates.

interrogated the reactivity of α -aryl vinyl azides **3**. As shown in Figure 3, these substrates exhibited migration selectivity that could be turned by the electronic nature of the arene substituent: specifically, difluorination was accompanied by 1,2-migration of either the azide or aryl group, leading to β -difluoroalkylazides **4** or α -difluoroalkylazides **5**, respectively (Figure 3A). Azides with strong electron-withdrawing groups (EWGs) on the aryl moiety (**3aa–3ad**) underwent the expected *gem*-difluorination and 1,2-azide migration at 40°C to afford the corresponding products **4aa–4ad** in moderate to good yields (62%–82%, Figure 3B; inferior yields were obtained at 25°C). Certain disubstituted aryl vinyl azides, including 3,5-bis-trifluoromethyl- (**3ae**) and 3,5-difluoroarenes (**3af**), also afforded the azide migration products (**4ae** and **4af**), whereas substrates bearing formyl, acetyl, or ester groups at the *p*- or *m*-positions of the aryl ring (**3ag–3aj**) afforded mixtures of competing 1,2-azide and 1,2-aryl migrations, in favor of the former (**4ag–4aj** versus **4ag'–4aj'**), and the structure of azide migration product **4ah** was unambiguously confirmed by single-crystal X-ray crystallography. Realizing that the electronic nature of the arene could be used to control the chemoselectivity of the migration, other substituted arenes were tested. A meta-chloro substituent (**3ak**) again gave rise to a mixture of the products of the 1,2-migration (**4ak** and **4ak'**); however, chloro, bromo, and fluoro substituents at the *p*-position of the aryl ring all exclusively delivered the *gem*-difluorinated products of 1,2-aryl migration (**4al–4an**, 72%–79%). Despite the inductive electron-withdrawing nature of these substituents, this selectivity likely arises from mesomeric stabilization of developing positive charge at the *para* position during arene migration. This switch in product selection allowed us to prepare a variety of previously inaccessible α -difluoroalkylazides **5**, thus providing an entry to novel α -difluoroalkylamines.

This switch in selectivity was conveniently extended to a range of α -aryl vinyl azides equipped with electron-donating substituents (EDGs) on the aryl moiety (Figure 3B). α -aryl vinyl azides featuring alkyl and alkoxy groups at any position on the aryl ring (**3ba–3bi**) were efficiently transformed into the corresponding 1,2-aryl migration products (α -difluoroalkylazides **5ba–5bi**) in good to excellent yields, and no 1,2-azide migration products were observed. Substrates with additional sp^2 substituents (2-naphthyl, phenyl, and 3-thienyl; **3bj–bl**) were also smoothly converted to equivalent products (**5bj–5bl**) in high yields, and the structure of **5bk** was confirmed by single-crystal X-ray crystallography. Overall, these results illustrate the connection between migratory aptitude and the arene substituent (EWG versus EDG).

The vinyl azides employed in this chemistry are readily derived by hydroazidation of the corresponding terminal alkynes,⁴⁹ and many tandem hydroazidation reactions of terminal alkynes have been established.^{50,51} Accordingly, we questioned whether it would be possible to access the difluoroalkylazide products directly from alkynes in a two-step, one-pot process consisting of hydroazidation, followed by *gem*-difluorination/1,2-azide migration. Using alkyne **6** as substrate, both steps indeed proceeded efficiently, giving *gem*-difluorinated alkyl azide **2aa** in 80% yield (Figure 4A). This sequenced approach offers a step economic and operationally simple method for the synthesis of *gem*-difluorinated alkyl azides.

The practicality of the process was readily tested by conducting a multigram scale synthesis of **2aa** (Figure 4B, 15 mmol scale, 2.7 g, 87%). **2aa** can be used to




Figure 3. Divergent Migration Pathways in the *gem*-difluorination of α -Aryl Vinyl Azides

synthesize various previously unknown *gem*-difluorinated compounds of potential interest as useful building blocks in medicinal chemistry: for instance, reduction of the azide group produced β -difluoroalkylamine **7** in excellent yield,⁴² whereas base-mediated 1,3-dipolar cycloaddition of **2aa** with malononitrile⁵² or dimethyl acetylenedicarboxylate (DMAD)⁵³ gave the trisubstituted 1,2,3-triazole derivatives **8** and **9** in 65% and 95% yields, respectively. 1,3-dipolar cycloaddition of **2aa** with benzoyl cyanide afforded 1,5-disubstituted tetrazole **10** in good yield,⁵⁴ whereas addition of an alkyl lithium afforded triazene **11** in 90% yield,⁵⁵ the practical dipolar cycloaddition could further promote their use in bioconjugation.^{12,56} These results clearly illustrate the potential of β -difluorinated alkyl azides to access nitrogen-containing β -difluorinated products, which could influence physical and biological properties such as conformation, lipophilicity, and metabolic stability.^{57–59} Otherwise, positron emission tomography (PET) and single photon emission computed tomography (SPECT) are subfields of nuclear imaging that emerged from the combined efforts of national laboratories, academia, and industries.^{60,61} Fluorine-18 (¹⁸F, $t_{1/2}$ = 109.7 min) is the most widely used nuclide for positron emission tomography (PET). Our methodology has quite a rapid reaction time, even occurring at speeds <1 min. Such a short reaction time should be essential for the efficient introduction of ¹⁸F-radiolabel into organic molecules.^[62]



A 1,2-azide migratory *gem*-difluorination starting from terminal alkynesB Gram scale synthesis and further reactions of the β -difluoroazides to medicinal chemistry intermediates

Figure 4. Tandem Hydroazidation/Difluorination, Gram Scale Synthesis, and Further Product Transformations

(A) *gem*-difluorination and 1,2-azide migration of terminal alkynes. Reaction conditions: **6** (0.5 mmol), TMSN_3 (1.0 mmol), Ag_2CO_3 (10% mol), in DMSO (2.0 mL) and H_2O (2.0 equiv) at 80°C for 90 min; then DCM (2 mL), $\text{Py}\cdot\text{HF}$ (2.5 mmol), PIDA (1.0 mmol), 25°C . PIDA = (Diacetoxyiodo)benzene; Py = pyridyl.

(B) Gram scale synthesis and further reactions of the β -difluoroalkylazides to access products relevant to medicinal chemistry applications. Reaction conditions: (a) **2aa** (0.5 mmol), PPh_3 (1.0 mmol), H_2O (200 μL) in THF (2.0 mL) at 50°C for 3 h. (b) **2aa** (0.5 mmol), malononitrile (0.6 mmol), K_2CO_3 (1.0 mmol) in DMSO at 45°C . (c) **2aa** (0.5 mmol), dimethyl acetylenedicarboxylate (0.6 mmol) in water (8 mL) at 70°C . (d) **2aa** (0.5 mmol) and benzoyl cyanide (1.2 mmol) at 120°C under Ar. (e) **2aa** (0.5 mmol), *n*-BuLi (0.6 mmol) in THF at -78°C under Ar. THF = tetrahydrofuran.

The mechanism of azide group migration is intriguing. To explore this, the *gem*-difluorination/1,2-azide migration was investigated with deuterium-labeled substrate $[\text{D}_2]$ -**1aa** (Figure 5A), which under the standard conditions produced difluorinated product $[\text{D}_2]$ -**2aa** in 95% yield with exclusive deuteriation at the carbon atom bearing the azide group. No indication of deuterium scrambling was observed by ^1H NMR spectroscopic analysis, suggesting the involvement of a 1,2-azide migration rather than an elimination process. A mechanism is proposed that involves initial 1,2-iodofluorination of the vinyl azide with $\text{PhIF}_2\cdot 2\text{HF}$ (generated *in situ*




Figure 5. Mechanistic Study and Proposal for the *gem*-difluorination/1,2-Azide Migration of Vinyl Azides

The energy profile was obtained by DFT calculations at the M062X/6-31G(d, p)/LANL2DZ level.

from reaction of PIDA with Py·HF, forming intermediate I (Figure 5B). The similar activation of different molecules of HF to PhIF₂ has been investigated by Zhou et al.,³³ and they found that the two-molecule activation model is the most favored. Thus, we directly use this activation model to study the following processes in this work. Subsequently, azide migration (promoted by the leaving group ability of iodobenzene) and concomitant fluorination appear to outcompete processes involving azide decomposition driven by loss of N₂. This 1,2-azide shift in intermediate I could occur either via a five-membered ring transition state (TS-1, path a) or a three-membered ring transition state (TS-2, path b). These transition states were studied with density functional theory (DFT) calculations at the M062X/6-31G(d,p)/LANL2DZ level. The calculations predict that path a (via TS-1, Figure 5B) presents a prohibitively high energy barrier (67.0 kcal/mol) relative to path b (TS-2, 15.1 kcal/mol), potentially because of (1) the torsional strain involved in the migration of the azide group in TS-1 ($\theta_1 = 161.8^\circ$) makes its structure more destabilized than that of TS-2 ($\theta_2 = 174.4^\circ$); (2) the stronger hydrogen-bond interactions between F⁻ and H as




Figure 6. NCI analysis for the two key transition states TS-1 and TS-2

The colors blue, green, and red represent the attraction, weak interaction, and steric effect, respectively, between the atoms.

well as the stronger electrostatic interactions between F^- and I^+ in TS-2 is more favored than those in TS-1, the electrostatic interactions between F^- and N_2^+ on the azide group was also found in TS-2, whereas it is absent in TS-1 (NCI analysis in Figure 6). As such, path b is proposed to be favored. The calculations also show that the second fluorination occurs exclusively by nucleophilic attack at the fluorinated carbon, and the azonium ion is the leaving group, completing 1,2-azide migration to β -difluorinated alkyl azide product **2ab**. We speculate that the avoidance of azide decomposition could be explained by the intramolecular delivery of the fluoride ion. Finally, in the case of α -aryl vinyl azides, the electronic character of the substituent on the aryl ring plays a pivotal role in the migration selectivity. Whereas electron-poor aromatics undergo equivalent azide migration, the ability of electron-rich arenes to form a phenonium ion outcompetes 1,2-azide migration and thus delivers α -difluorinated alkyl azides.

Conclusions

An unprecedented 1,2-azide migration has been developed in the *gem*-difluorination of the easily available α -vinyl azides with *in situ*-generated $PhIF_2 \cdot HF$, providing efficient access to a variety of previously inaccessible, synthetically useful β -difluorinated azides. The azide functionality in the products can easily undergo further transformations to afford a variety of *gem*-difluorinated nitrogen-containing molecules of potential interest in medicinal chemistry. The azide migration could lead to those organic azides that are inaccessible by conventional methods; therefore, the discovery described here represents a conceptual advance in azide preparation and has opened up an avenue to the azide migration highly desired by synthetic chemists.

EXPERIMENTAL PROCEDURES

Full experimental procedures are provided in the [Supplemental Information](#).

DATA AND CODE AVAILABILITY

The X-ray crystallographic coordinates for structures reported in this article have been deposited at the Cambridge Crystallographic Data Center (4ah^{''}: 1960403, 5bk: 1893371). These data could be obtained free of charge from The Cambridge Crystallographic Data Center via <https://www.ccdc.cam.ac.uk/structures/>.

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.chempr.2019.12.004>.

ACKNOWLEDGMENTS

Financial support by NSFC (21871043, 21961130376), Department of Science and Technology of Jilin Province (20180101185JC and 20190701012GH), and the Fundamental Research Funds for the Central Universities (2412019ZD001). E.A.A. thanks the EPSRC for support (EP/M019195/1 and EP/S013172/1). X.B. and E.A.A. thank Royal Society for a Newton Advanced Fellowship.

AUTHOR CONTRIBUTIONS

Y.N. and P.S. performed the experiments and the mechanistic studies. Y.N., G.Z., E.A.A., and X.B. conceived the concept, designed the project, analyzed the data, and prepared this manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: June 13, 2019

Revised: September 8, 2019

Accepted: December 5, 2019

Published: December 30, 2019

REFERENCES

- Hili, R., and Yudin, A.K. (2006). Making carbon-nitrogen bonds in biological and chemical synthesis. *Nat. Chem. Biol.* **2**, 284–287.
- Ruiz-Castillo, P., and Buchwald, S.L. (2016). Applications of palladium-catalyzed C–N cross-coupling reactions. *Chem. Rev.* **116**, 12564–12649.
- Griess, P. (1864). On a new class of compounds in which nitrogen is substituted for hydrogen. *Proc. R. Soc. London* **13**, 375–384.
- Chiba, S. (2012). Application of organic azides for the synthesis of nitrogen-containing molecules. *Synlett* **2012**, 21–44.
- Shin, K., Kim, H., and Chang, S. (2015). Transition-metal-catalyzed C–N bond forming reactions using organic azides as the nitrogen source: a journey for the mild and versatile C–H amination. *Acc. Chem. Res.* **48**, 1040–1052.
- Fu, J., Zanon, G., Anderson, E.A., and Bi, X. (2017). α -Substituted vinyl azides: an emerging functionalized alkene. *Chem. Soc. Rev.* **46**, 7208–7228.
- Hayashi, H., Kaga, A., and Chiba, S. (2017). Application of vinyl azides in chemical synthesis: a recent update. *J. Org. Chem.* **82**, 11981–11989.
- Driver, T.G. (2013). C–H bond functionalization: an aminated reaction. *Nat. Chem.* **5**, 736–738.
- Sharma, A., and Hartwig, J.F. (2015). Metal-catalyzed azidation of tertiary C–H bonds suitable for late-stage functionalization. *Nature* **517**, 600–604.
- Lang, S., and Murphy, J.A. (2006). Azide rearrangements in electron-deficient systems. *Chem. Soc. Rev.* **35**, 146–156.
- Schilling, C.I., Jung, N., Biskup, M., Schepers, U., and Bräse, S. (2011). Bioconjugation via azide–staudinger ligation: an overview. *Chem. Soc. Rev.* **40**, 4840–4871.
- Kolb, H.C., Finn, M.G., and Sharpless, K.B. (2001). Click chemistry: diverse chemical function from a few good reactions. *Angew. Chem. Int. Ed.* **40**, 2004–2021.
- Bräse, S., Gil, C., Knepper, K., and Zimmermann, V. (2005). Organic azides: an exploding diversity of a unique class of compounds. *Angew. Chem. Int. Ed.* **44**, 5188–5240.
- S. Bräse, and K. Banert, eds. (2010). *Organic Azides: Syntheses and Applications* (John Wiley & Sons).
- L'abbe, G. (1969). Decomposition and addition reactions of organic azides. *Chem. Rev.* **69**, 345–363.
- Aguila, M.J.B., Badiei, Y.M., and Warren, T.H. (2013). Mechanistic insights into C–H amination via dicopper nitrenes. *J. Am. Chem. Soc.* **135**, 9399–9406.
- G. Soderberg, B.C.G. (2000). Synthesis of heterocycles via intramolecular annulation of nitrene intermediates. *Curr. Org. Chem.* **4**, 727–764.
- Uchida, T., and Katsuki, T. (2014). Asymmetric nitrene transfer reactions: sulfimidation, aziridination and C–H amination using azide compounds as nitrene precursors. *Chem. Rec.* **14**, 117–129.
- López, E., and López, L.A. (2017). Synthesis of functionalized cyclopentene derivatives from vinyl diazo compounds and vinyl azides through sequential copper-promoted [3+2] cycloaddition/azide rearrangement. *Angew. Chem. Int. Ed.* **56**, 5121–5124.
- Porter, M.R., Shaker, R.M., Calcanas, C., and Topczewski, J.J. (2018). Stereoselective dynamic cyclization of allylic azides: synthesis of tetralins, chromanes, and tetrahydroquinolines. *J. Am. Chem. Soc.* **140**, 1211–1214.
- Liu, R., Gutierrez, O., Tantillo, D.J., and Aubé, J. (2012). Stereocontrol in a combined allylic azide rearrangement and intramolecular Schmidt reaction. *J. Am. Chem. Soc.* **134**, 6528–6531.
- Ott, A.A., Goshey, C.S., and Topczewski, J.J. (2017). Dynamic kinetic resolution of allylic azides via asymmetric dihydroxylation. *J. Am. Chem. Soc.* **139**, 7737–7740.
- Packard, M.H., Cox, J.H., Suding, V.P., and Topczewski, J.J. (2017). The effect of proximal functionality on the allylic azide equilibrium. *Eur. J. Org. Chem.* **2017**, 6365–6368.
- Topczewski, J.J., and Porter, M.R. (2018). “It’s a (kinetic) trap!” - selectively differentiating allylic azide isomers. *Synlett* **29**, 1537–1542.
- Smith, G.R., Brenneman, D.E., Zhang, Y., Du, Y., and Reitz, A.B. (2014). Small-molecule anticonvulsant agents with potent *in vitro* neuroprotection and favorable drug-like properties. *J. Mol. Neurosci.* **52**, 446–458.
- Cox, C.D., Coleman, P.J., Breslin, M.J., Whitman, D.B., Garbaccio, R.M., Fraley, M.E., Buser, C.A., Walsh, E.S., Hamilton, K., Schaber, M.D., et al. (2008). Kinesin spindle protein (KSP) inhibitors. 9. Discovery of (2S)-4-(2,5-difluorophenyl)-n-[(3R,4S)-3-fluoro-1-methylpiperidin-4-yl]-2-(hydroxymethyl)-N-methyl-2-phenyl-2,5-dihydro-1H-pyrrole-1-carboxamide (MK-0731) for the treatment of taxane-refractory cancer. *J. Med. Chem.* **51**, 4239–4252.
- Welch, J.T., and Eswarakrishnan, S. (1991). *Fluorine in Bioorganic Chemistry* (John Wiley & Sons).

28. R.E. Banks, B.E. Smart, and J.C. Tatlow, eds. (1994). *Organofluorine Chemistry: Principles and Commercial Applications* (Springer US).
29. Müller, K., Faeh, C., and Diederich, F. (2007). Fluorine in pharmaceuticals: looking beyond intuition. *Science* 317, 1881–1886.
30. Meanwell, N.A. (2018). Fluorine and fluorinated motifs in the design and application of bioisosteres for drug design. *J. Med. Chem.* 61, 5822–5880.
31. Seebach, D., Dubost, E., Mathad, R., Jaun, B., Limbach, M., Löweneck, M., Flögel, O., Gardiner, J., Capone, S., Beck, A., et al. (2008). New open-chain and cyclic tetrapeptides, consisting of α -, β^2 -, and β^3 -amino acid residues, as somatostatin mimics- a survey. *Helv. Chim. Acta* 91, 1736–1786.
32. Banik, S.M., Medley, J.W., and Jacobsen, E.N. (2016). Catalytic, asymmetric difluorination of alkenes to generate difluoromethylated stereocenters. *Science* 353, 51–54.
33. Zhou, B., Haj, M.K., Jacobsen, E.N., Houk, K.N., and Xue, X.S. (2018). Mechanism and origins of chemo- and stereoselectivities of aryl iodide-catalyzed asymmetric difluorinations of β -substituted styrenes. *J. Am. Chem. Soc.* 140, 15206–15218.
34. Ilchenko, N.O., Tasch, B.O.A., and Szabó, K.J. (2014). Mild silver-mediated geminal difluorination of styrenes using an air- and moisture-stable fluoroiodane reagent. *Angew. Chem. Int. Ed.* 53, 12897–12901.
35. Zhao, Z., Racicot, L., and Murphy, G.K. (2017). Fluorinative rearrangements of substituted phenylallenes mediated by (difluoroiodo) toluene: synthesis of α -(difluoromethyl) styrenes. *Angew. Chem. Int. Ed.* 56, 11620–11623.
36. Lv, W.X., Li, Q., Li, J.L., Li, Z., Lin, E., Tan, D.H., Cai, Y.H., Fan, W.X., and Wang, H. (2018). *gem*-difluorination of alkenyl *n*-methyliminodiacetyl boronates: synthesis of α - and β -difluorinated alkylborons. *Angew. Chem. Int. Ed.* 57, 16544–16548.
37. Scheidt, F., Neufeld, J., Schäfer, M., Thiehoff, C., and Gilmour, R. (2018). Catalytic geminal difluorination of styrenes for the construction of fluorine-rich bioisosteres. *Org. Lett.* 20, 8073–8076.
38. Kitamura, T., Yoshida, K., Mizuno, S., Miyake, A., and Oyamada, J. (2018). Difluorination of functionalized aromatic olefins using hypervalent iodine/HF reagents. *J. Org. Chem.* 83, 14853–14860.
39. Meanwell, N.A. (2011). Synopsis of some recent tactical application of bioisosteres in drug design. *J. Med. Chem.* 54, 2529–2591.
40. Mennie, K.M., Banik, S.M., Reichert, E.C., and Jacobsen, E.N. (2018). Catalytic diastereo- and enantioselective fluoroamination of alkenes. *J. Am. Chem. Soc.* 140, 4797–4802.
41. Fu, N., Sauer, G.S., Saha, A., Loo, A., and Lin, S. (2017). Metal-catalyzed electrochemical diazidation of alkenes. *Science* 357, 575–579.
42. Yuan, Y.A., Lu, D.F., Chen, Y.R., and Xu, H. (2016). Iron-catalyzed direct diazidation for a broad range of olefins. *Angew. Chem. Int. Ed.* 55, 534–538.
43. Gao, B., Zhao, Y., and Hu, J. (2015). AgF-mediated fluorinative cross-coupling of two olefins: facile access to α -CF₃ alkenes and β -CF₃ ketones. *Angew. Chem. Int. Ed.* 54, 638–642.
44. Xu, P., Wang, F., Fan, G., Xu, X., and Tang, P. (2017). Hypervalent iodine(III)-mediated oxidative fluorination of alkylsilanes by fluoride ions. *Angew. Chem. Int. Ed.* 56, 1101–1104.
45. Huang, X., Liu, W., Hooker, J.M., and Groves, J.T. (2015). Targeted fluorination with the fluoride ion by manganese-catalyzed decarboxylation. *Angew. Chem. Int. Ed.* 54, 5241–5245.
46. Qin, A., Lam, J.W.Y., and Tang, B.Z. (2010). Click polymerization. *Chem. Soc. Rev.* 39, 2522–2544.
47. Johnson, J.A., Finn, M.G., Koberstein, J.T., and Turro, N.J. (2008). Construction of linear polymers, dendrimers, networks, and other polymeric architectures by copper-catalyzed azide-alkyne cycloaddition “click” chemistry. *Macromol. Rapid Commun.* 29, 1052–1072.
48. Svenson, S. (2015). The dendrimer paradox—high medical expectations but poor clinical translation. *Chem. Soc. Rev.* 44, 4131–4144.
49. Liu, Z., Liao, P., and Bi, X. (2014). General silver-catalyzed hydroazidation of terminal alkynes by combining TMS-N₃ and H₂O: synthesis of vinyl azides. *Org. Lett.* 16, 3668–3671.
50. Ning, Y., Ji, Q., Liao, P., Anderson, E.A., and Bi, X. (2017). Silver-catalyzed stereoselective aminosulfonylation of alkynes. *Angew. Chem. Int. Ed.* 56, 13805–13808.
51. Liu, B., Ning, Y., Virelli, M., Zanoni, G., Anderson, E.A., and Bi, X. (2019). Direct transformation of terminal alkynes into amidines by a silver-catalyzed four-component reaction. *J. Am. Chem. Soc.* 141, 1593–1598.
52. Tome, A.C. (2004). In *Science of synthesis: Houben-Weyl methods of molecular transformations*, 13, R.C. Storr and T.L. Gilchrist, eds. (Thieme), pp. 415–601.
53. Sun, X., Li, X., Song, S., Zhu, Y., Liang, Y.F., and Jiao, N. (2015). Mn-catalyzed highly efficient aerobic oxidative hydroxyazidation of olefins: a direct approach to β -azido alcohols. *J. Am. Chem. Soc.* 137, 6059–6066.
54. Demko, Z.P., and Sharpless, K.B. (2002). A click chemistry approach to tetrazoles by Huisgen 1,3-dipolar cycloaddition: synthesis of 5-acyltetrazoles from azides and acyl cyanides. *Angew. Chem. Int. Ed.* 41, 2113–2116.
55. Sieh, D.H., Wilbur, D.J., and Michejda, C.J. (1980). Preparation of trialkyltriazenes. A comparison of the nitrogen-nitrogen bond rotation in trialkyltriazenes and aryltrialkyltriazenes by variable temperature carbon-13 NMR. *J. Am. Chem. Soc.* 102, 3883–3887.
56. Thirumurugan, P., Matosiuk, D., and Jozwiak, K. (2013). Click chemistry for drug development and diverse chemical–biology applications. *Chem. Rev.* 113, 4905–4979.
57. Purser, S., Moore, P.R., Swallow, S., and Gouverneur, V. (2008). Fluorine in medicinal chemistry. *Chem. Soc. Rev.* 37, 320–330.
58. Wang, J., Sánchez-Roselló, M., Aceña, J.L., del Pozo, C., Sorochinsky, A.E., Fustero, S., Soloshonok, V.A., and Liu, H. (2014). Fluorine in pharmaceutical industry: fluorine-containing drugs introduced to the market in the last decade (2001–2011). *Chem. Rev.* 114, 2432–2506.
59. Gillis, E.P., Eastman, K.J., Hill, M.D., Donnelly, D.J., and Meanwell, N.A. (2015). Applications of fluorine in medicinal chemistry. *J. Med. Chem.* 58, 8315–8359.
60. Preshlock, S., Tredwell, M., and Gouverneur, V. (2016). (18)F-labeling of arenes and heteroarenes for applications in positron emission tomography. *Chem. Rev.* 116, 719–766.
61. Constanze, N.N., Jacob, M., and Hooker, T.R. (2016). Concerted nucleophilic aromatic substitution with ¹⁹F⁻ and ¹⁸F⁻. *Nature* 534, 369–373.
62. Miguel, A.C.G., Patrik, N., Antonio, B.G., Denise, N.M., Gunnar, A., Magnus, S., and Kálmán, J.S. (2018). [¹⁸F]fluoro-benziodoxole: a no-carrier-added electrophilic fluorinating reagent. Rapid, simple radiosynthesis, purification and application for fluorine-18 labelling. *Chem. Commun. (Camb.)* 54, 4286–4289.