


# Photocatalyzed Site-selective C-H to C-C Conversion of Aliphatic Nitriles

Keiichi Yamada,<sup>†</sup> Megumi Okada,<sup>†</sup> Takahide Fukuyama,<sup>†</sup> Davide Ravelli,<sup>‡</sup> Maurizio Fagnoni,<sup>\*,‡</sup> and Ilhyong Ryu<sup>\*,†</sup>

<sup>†</sup>Department of Chemistry, Graduate School of Science, Osaka Prefecture University, Sakai, Osaka 599-8531, Japan.

<sup>‡</sup>PhotoGreen Lab, Department of Chemistry, University of Pavia, Viale Taramelli 12, 27100 Pavia, Italy.


Supporting Information Placeholder



**ABSTRACT:**  $\beta$ - or  $\gamma$ -Site-selective C-H alkylation of aliphatic nitriles has been achieved using a decatungstate salt as the photocatalyst. The observed site-selectivity was justified by a radical polar effect in transition states for hydrogen abstraction.


The site-selective functionalization of alkanes is currently one of the most important challenges in organic chemistry.<sup>1</sup> Aliphatic nitriles are potentially useful compounds, since they can be easily converted to a variety of carbonyl-functionalized compounds, such as carboxylic acids,<sup>2</sup> aldehydes,<sup>3</sup> esters,<sup>4</sup> amines,<sup>5</sup> and amides.<sup>6</sup> We theorized that functionalization of the C-H bonds in aliphatic nitriles would significantly widen the scope of accessible nitriles. Remarkably, site-selective C-H/C-C conversion in aliphatic nitriles is not readily achieved in any position other than  $\alpha$ -to the nitrile.<sup>7,8</sup> Recently, we reported that cyclopentanones could be regioselectively functionalized at the  $\beta$ -position using tetrabutylammonium decatungstate (TBADT)<sup>9</sup> as a photocatalyst, leading to highly selective C-H alkylation and acylation.<sup>10</sup> The polar effect<sup>11</sup> exerted in the  $S_{\text{H}}2$  transition state of hydrogen abstraction by the excited decatungstate anion was invoked to explain the lack of reactivity at the usually activated  $\alpha$ -position. In the present study, we report that this photocatalytic approach was highly effective for the  $\beta$ - and  $\gamma$ -site-selective alkylation of aliphatic nitriles by alkenes (Scheme 1).

## Scheme 1. Site-selective Alkylation of Aliphatic Nitriles Presented in This Work



As a model reaction, we chose to study the alkylation of butyronitrile (**1a**) by dimethyl maleate (**2a**; Scheme 2). Thus, an acetonitrile (1 mL) solution of **1a** (20 equiv), **2a** (0.5 mmol), and TBADT (4 mol%) was irradiated using a Xe lamp (300 W) through a Pyrex tube for 24 h. Under these conditions,  $\beta$ -substituted butyronitrile **3a** was obtained in 69% isolated yield as a 51:49 mixture of diastereoisomers (Scheme 2, eq 1), while  $\alpha$ -substituted butyronitrile was not observed, and  $\gamma$ -substituted butyronitrile **4a** was formed in only 7% yield. The complete avoidance of  $\alpha$ -C-H activation can be accounted for by an unfavorable polar transition state for hydrogen abstraction  $\alpha$  to the nitrile group (vide infra, Scheme 5, TS-A (R = H)). The preference for  $\beta$ -methylene activation rather than  $\gamma$ -methyl activation in **1a** reflects the weaker bond strength of the methylene C-H bond. Next, we examined the alkylation of valeronitrile (**1b**), which has both  $\beta$ - and  $\gamma$ -methylene groups, with **2a** (Scheme 2, eq 2). The reaction yielded  $\gamma$ -substituted nitrile **4b** in a noteworthy preference to  $\beta$ -substituted nitrile **3b** ( $\gamma/\beta = 83/14$ ). In this reaction, products derived from C-H cleavage at the methyl position yielded only a trace amount of **5b**. The reaction of isovaleronitrile (**1c**), bearing a methine carbon at the  $\beta$ -position, with **2a** gave the corresponding  $\beta$ -functionalized nitrile **3c** in 79% yield with complete site-selectivity (Scheme 2, eq 3).

## Scheme 2. Site-selectivity Observed in the TBADT Photocatalyzed C-H Alkylation of Butyronitrile (**1a**), Valeronitrile (**1b**), and Isovaleronitrile (**1c**) by Dimethyl Maleate (**2a**)



After obtaining promising results for the site-selective C-H functionalization of **1a**, **1b**, and **1c**, we next sought to assess the selectivity of a variety of aliphatic nitriles **1** in the reaction with electron-deficient alkenes **2**. The results are summarized in Table 1. In a similar manner, the reactions of **1c** with *t*-butyl acrylate (**2b**), methyl vinyl ketone (**2c**), acrylonitrile (**2d**), acrylamide (**2e**), and phenyl vinyl sulfone (**2f**) all proceeded selectively at the  $\beta$ -position to give the corresponding  $\beta$ -alkylated nitriles **3d-h** in good yields (entries 1-5). The reaction of propionitrile (**1d**) with **2a** occurred selectively at the  $\beta$ -methyl, but it was very sluggish, giving product **3i** in a low yield after 48 h (entry 6). The alkylation of cyclopropyl cyanide (**1e**) with **2a** did not proceed (entry 7). The reaction of isocaproitrile (**1f**) with **2a** proceeded selectively at the  $\gamma$ -methine to give **3k** in 79% yield (entry 8). The reaction of **1f** with alkenes **2b**, **2c**, **2d**, and **2g** also gave the corresponding  $\gamma$ -alkylation products **3l**, **3m**, **3n**, and **3o**, respectively, in good yields (entries 9-12). Notably, the formation of compound **3o** was likewise successfully when the irradiation source was solar simulated light. The reaction of dinitriles was also examined. While the reaction of adiponitrile (**1g**) with **2a** proceeded selectively at the  $\beta$ -methylene carbon to give the  $\beta$ -alkylated product **3p** (entry 13), the reaction of glutaronitrile (**1h**) with **2a** did not proceed (entry 14). Dinitrile **1i**, which has six different C-H bonds including both nucleophilic and electrophilic methine groups, reacted with **2a** selectively at the only nucleophilic methine carbon to give **3r** in 52% yield (entry 15). The direct sunlight-induced alkylation of aliphatic nitrile **1f** was also successful, and gave  $\gamma$ -alkylated nitrile **3k** in a 62% yield (entry 16).


Table 1. Site-selective C-H Alkylation by Aliphatic Nitriles **1** Using TBADT as a Photocatalyst<sup>a</sup>

Entry	Nitrile <b>1</b>	Alkene <b>2</b>	Product <b>3</b>	Yield <sup>b</sup>
1	<b>1c</b>	<b>2b</b>	<b>3d</b>	82%
2	<b>1c</b>	<b>2c</b>	<b>3e</b>	86%
3	<b>1c</b>	<b>2d</b>	<b>3f</b>	73%
4 <sup>c</sup>	<b>1c</b>	<b>2e</b>	<b>3g</b>	68%
5	<b>1c</b>	<b>2f</b>	<b>3h</b>	83%
6 <sup>d</sup>	<b>1d</b>	<b>2a</b>	<b>3i</b>	31%
7	<b>1e</b>	<b>2a</b>	<b>3j</b>	n.d. <sup>e</sup>
8 <sup>f</sup>	<b>1f</b>	<b>2a</b>	<b>3k</b>	79%
9 <sup>f</sup>	<b>1f</b>	<b>2b</b>	<b>3l</b>	75%
10 <sup>f</sup>	<b>1f</b>	<b>2c</b>	<b>3m</b>	60%
11 <sup>f</sup>	<b>1f</b>	<b>2d</b>	<b>3n</b>	73%
12 <sup>g</sup>	<b>1f</b>	<b>2g</b>	<b>3o</b>	54% (only <i>endo</i> )
13	<b>1g</b>	<b>2a</b>	<b>3p</b>	69%
14	<b>1h</b>	<b>2a</b>	<b>3q</b>	n.d. <sup>e</sup>
15	<b>1i</b>	<b>2a</b>	<b>3r</b>	52%
16 <sup>h</sup>	<b>1f</b>	<b>2a</b>	<b>3k</b>	62%


<sup>a</sup> **1** (10 mmol), **2** (0.5 mmol), TBADT (4 mol%), acetonitrile (1 mL), irradiation by 300 W Xe lamp for 24 h. <sup>b</sup> Yields of product isolated after flash chromatography on SiO<sub>2</sub>. If necessary, further purification was made by preparative HPLC. <sup>c</sup> 1 mL of acetonitrile/acetone (1/1) was used as the solvent. <sup>d</sup> 48 h. <sup>e</sup> Not detected. <sup>f</sup> Trace amount of  $\beta$ -alkylated nitrile was detected by <sup>13</sup>C NMR of the crude reaction mixture. <sup>g</sup> **1** (10 mmol), **2** (0.5 mmol), TBADT (4 mol%), acetonitrile (5 mL), irradiation by a SolarBox equipped with a 1.5 kW Xe lamp (500 Wm<sup>-2</sup>) for 20 h. <sup>h</sup> Sunlight exposure for 56 h (7 days) in Osaka (autumn).

To determine whether the key alkyl radical species would react with CO,<sup>12</sup> we examined a TBADT-catalyzed three-component coupling reaction comprising butyronitrile (**1a**), CO, and dimethyl maleate (**2a**). The reaction gave a mixture of  $\beta$ -acylated product **6a** and  $\gamma$ -acylated product **7a** in a 75/25 ratio under high CO pressures (Scheme 3). The decline of  $\beta$ -selectivity compared with alkylation using **2a** ( $\beta/\gamma = 91/9$ ) may be ascribed to the stability of acyl radicals towards backward decarbonylation, since loss of CO from a secondary acyl radical is faster than that from a primary acyl radical.<sup>12</sup>

**Scheme 3. Three-component,  $\beta$ -Selective C-H Acylation of Butyronitrile (**1a**)**



To gain insight into the mechanism, we carried out laser flash photolysis experiments. The reactive excited state of decatungstate shows a strong absorption in the red part of the visible spectrum and can be revealed at  $\lambda = 780$  nm.<sup>13</sup> In the presence of hydrogen donors, a long-lived signal can be likewise observed and has been attributed to the reduced form of decatungstate ( $H^+[W_{10}O_{32}]^{5-}$ ) that absorbs in the same region.<sup>9e,13</sup> The signal decay profiles resulting after laser excitation of the decatungstate anion were registered in acetonitrile, propionitrile and butyronitrile. The signal lifetime was longer in acetonitrile (62 ns) compared to propionitrile (46 ns) and butyronitrile (39 ns; Figure 1 and Table S1). An opposite trend was observed for the long-lived signal, the intensity of which increased when shifting from acetonitrile to butyronitrile. Both spectral features are consistent with a more efficient hydrogen atom abstraction by excited decatungstate from butyronitrile than from acetonitrile, with propionitrile in between (see SI for further details).




**Figure 1.** Nanosecond decays observed at 780 nm after 355 nm laser excitation of a  $2 \times 10^{-4}$  M TBADT solution in: acetonitrile (red line), propionitrile (blue line) and butyronitrile (black line).


Scheme 4 illustrates a plausible reaction mechanism for the  $\beta$ -site-selective C-H alkylation of butyronitrile (**1a**) with dimethyl maleate (**2a**). The excited decatungstate anion abstracts a hydrogen from the  $\beta$ -C-H bond of **1a** to form the  $\beta$ -cyano radical **A**, which then combines with **2a** to form the adduct radical **B**. Back-hydrogen atom transfer from the re-

duced decatungstate anion gives the alkylated product **3a**, and regenerates the starting decatungstate anion. The reluctant  $\alpha$ -C-H functionalization of **1a** is rationalized by an unfavorable polar effect in the transition state **TS-A** ( $R = H$ ; Scheme 5), in which a partial positive charge on the  $\alpha$ -carbon is destabilized by the adjacent electronegative cyano group. The preference of  $\beta$ -methylene C-H to  $\gamma$ -methyl C-H may reasonably be ascribed to a weaker methylene C-H than the methyl C-H bond.<sup>14</sup> In the alkylation of valeronitrile (**1b**), a preference for  $\gamma$ -methylene with respect to  $\beta$ -methylene functionalization was observed. This suggests that the inductive effect of a cyano group is persistent in **TS-B** for  $\beta$ -C-H cleavage but not in **TS-C** for  $\gamma$ -C-H cleavage, and in this regard it should be noted that a similar tendency for  $\gamma$ -functionalization was recognized in the radical chlorination of valeronitrile.<sup>15,16,17</sup> This assumption could explain the failure of the reaction of **1h**, in which double destabilization of the polar **TS-D** by two  $\beta$ -cyano groups is present. In the successful case of adiponitrile (**1g**), the second cyano group that was located at the more remote  $\gamma$ -position exerted only a small effect, as shown in **TS-E**.

**Scheme 4. Proposed Reaction Mechanism for the Photocatalyzed  $\beta$ -C-H Alkylation of Butyronitrile (**1a**) with Dimethyl Maleate (**2a**)**



**Scheme 5. Possible Polar  $S_H2$  Transition States**



The present study used TBADT as a photocatalyst for the site-selective C-H to C-C conversion in aliphatic nitriles **1** with electron deficient alkenes **2**. In all cases examined, no alkylation was observed at the weakest C-H bonds  $\alpha$  to the cyano group, resulting in high  $\beta$ -site-selectivity for alkylation of butyronitrile. When aliphatic nitriles bearing both  $\beta$ - and  $\gamma$ -methylene hydrogens are reacted,  $\gamma$ -functionalization was preferred. When both methine and methylene groups were available, selective methine C-H cleavage was observed. These site-selective C-H cleavages in aliphatic nitriles led to a

new protocol for the synthesis of higher nitriles from lower nitriles. This *polar* radical approach could become a powerful and general tool for the site-selective C-H functionalization. The procedure is very straightforward and can be easily carried out by solar light irradiation. We are now applying this strategy to other systems.

## ASSOCIATED CONTENT

### Supporting Information

This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Author

Ilhyong Ryu

Department of Chemistry, Graduate School of Science, Osaka Prefecture University, Sakai, Osaka 599-8531, Japan. Fax: (+81) 72 254 9695; Tel: (+81) 72 254 9695; E-mail: [ryu@c.s.osakafu-u.ac.jp](mailto:ryu@c.s.osakafu-u.ac.jp)

Maurizio Fagnoni

PhotoGreen Lab, Department of Chemistry, University of Pavia, viale Taramelli 12, 27100 Pavia, Italy. Fax: (+39) 0382 987323; Tel: (+39) 0382 987198; E-mail: [fagnoni@unipv.it](mailto:fagnoni@unipv.it)

### Author Contributions

The manuscript was written through contributions of all authors. / All authors have given approval to the final version of the manuscript. / ‡These authors contributed equally. (match statement to author names with a symbol)

## ACKNOWLEDGMENT

This work was supported by a Grant-in-Aid for Scientific Research from the MEXT and the JSPS. We thank Miss Gökçe Türkkan (Ankara University) for preliminary experiments.

## REFERENCES

(1) For selected references on site-selective C-H functionalization, see: (a) Chen, H.; Schlecht, S.; Semple, T. C.; Hawtwig, J. F. *Science* **2000**, *287*, 1995. (b) Chen, M. S.; White, M. C. *Science* **2007**, *318*, 783. (c) Brodsky, B. H.; Bois, J. D. *J. Am. Chem. Soc.* **2005**, *127*, 15391. (d) Jeremy, K. C.; Richter, M.; Baran, P. S. *J. Am. Chem. Soc.* **2008**, *130*, 7247. (e) Chen, K.; Baran, P. S. *Nature* **2009**, *459*, 824. (f) Chen, M. S.; White, M. C. *Science* **2010**, *327*, 566. (g) Bigi, M. A.; Reed, S. A.; White, M. C. *J. Am. Chem. Soc.* **2012**, *134*, 9721. (h) Simmons, E. M.; Hartwig, J. F. *Nature* **2012**, *483*, 70. (i) Liu, W.; Huang, X.; Cheng, M. -J.; Nielsen, R. J.; Goddard III, W. A.; Groves, J. T. *Science* **2012**, *337*, 1322. (j) Bess, E. N.; DeLuca, R. J.; Tindall, D. J.; Oderinde, M. S.; Roizen, J. L.; Du Bois, J.; Sigman, M. S. *J. Am. Chem. Soc.* **2014**, *136*, 5783. (k) Kang, T.; Kim, Y.; Lee, D.; Wang, Z.; Chang, S. *J. Am. Chem. Soc.* **2014**, *136*, 4141. (l) Xia, J. -B.; Ma, Y.; Chen, C. *Org. Chem. Front.* **2014**, *1*, 468. (m) Li, B.; Driess, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2014**, *136*, 6586. (n) Kang, T.; Kim, H.; Kim, J. G.; Chang, S. *Chem. Commun.* **2014**, *50*, 12073. (o) Amaoka, Y.; Nagatomo, M.; Watanabe, M.; Tao, K.; Kamijo, S.; Inoue, M. *Chem. Sci.* **2014**, *5*, 4339. (p) Schmidt, V. A.; Quinn, R. K.; Brusoe, A. T.; Alexanian, E. J. *J. Am. Chem. Soc.* **2014**, *136*, 14389. (q) Alderson, J. M.; Phelps, A. M.; Scamp, R. J.; Dolan, N. S.; Schomaker, J. M. *J. Am. Chem. Soc.* **2014**, *136*, 16720. For recent reviews on transition-metal-catalyzed site-selective C-H functionalization, see: (r) Daugulis, O.; Do, H. -Q.; Shabashov, D. *Acc. Chem. Res.* **2009**, *42*, 1074. (s) Chen, X.; Engle, K. M.; Wang, D. -H.; Yu, J. -Q. *Angew. Chem., Int. Ed.* **2009**, *48*, 5094. (t) Jazzar, R.; Hitce, J.; Renaudat, A.; Sofack-Kreutzer, J.; Baudoin, O.

*Chem. Eur. J.* **2010**, *16*, 2654. (u) Newhouse, T.; Baran, P. S.; *Angew. Chem., Int. Ed.* **2011**, *50*, 3362. (v) Yan, G.; Borah, A. J. *Org. Chem. Front.* **2014**, *1*, 838. (w) Huang, Z.; Dong, G. *Tetrahedron Lett.* **2014**, *55*, 5869.

(2) Roy, A.; Roberts, F. G.; Wilderman, P. R.; Zhou, K.; Peter, R. J.; Coates, R. M. *J. Am. Chem. Soc.* **2007**, *129*, 12453.

(3) Snyder, S.; Treitler, D. S.; Brucks, A. P.; Sattler, W. *J. Am. Chem. Soc.* **2011**, *133*, 15898.

(4) Naota, T.; Shichijo, Y.; Murahashi, S. *J. Chem. Soc., Chem. Commun.* **1994**, 1359.

(5) (a) Liu, S.; Yang, Y.; Zhen, X.; Li, J.; He, H.; Feng, J.; Whiting, A. *Org. Biomol. Chem.* **2012**, *10*, 663. (b) Gunanathan, C.; Holscher, M.; Leitner, W. *Eur. J. Inorg. Chem.* **2011**, 3381.

(6) (a) Saito, E.; Naota, T.; Murahashi, S. *J. Am. Chem. Soc.* **1986**, *108*, 7846. (b) Naota, T.; Saito, E.; Murahashi, S. *J. Org. Chem.* **1992**, *57*, 2521. (c) Goto, A.; Endo, K.; Saito, S. *Angew. Chem., Int. Ed.* **2008**, *47*, 3607.

(7) For recent work of  $\alpha$ -functionalization of aliphatic nitriles, see: (a) Maruoka, K.; Kitamura, M.; Wang, X. *J. Am. Chem. Soc.* **2007**, *129*, 1038. (b) Concellon, J. M.; Rodeiguez-Solla, H.; Simal, C.; Santos, D.; Paz, N. R. *Org. Lett.* **2008**, *10*, 4549. (c) Wagener, K. B.; Wei, Y.; Inci, B.; Rojas, G. *J. Am. Chem. Soc.* **2007**, *129*, 17376. (d) Fleming, F. F.; Heppekausen, J.; Bernhardt, S.; Duez, K. P. *Org. Lett.* **2011**, *13*, 1690. (e) Mycka, R. J.; Eckenhoff, W. T.; Steward, O. W.; Barefoot, N. Z.; Fleming, F. F. *Tetrahedron* **2013**, *69*, 366. (f) Nambo, M.; Yar, M.; Smith, J. D.; Crudden, C. M. *Org. Lett.* **2015**, *17*, 50. Also, see a review: Bellina, F.; Rossi, R. *Chem. Rev.* **2010**, *110*, 1082.

(8) For aerobic oxidation of aliphatic nitriles leading to hydroperoxy-nitriles using decatungstate cluster as a photocatalyst, see: Ermolenko, L. P.; Giannotti, C. *J. Chem. Soc., Perkin Trans. 2* **1996**, 1205.

(9) For recent work on TBADT-catalyzed reaction, see: (a) Ryu, I.; Tani, A.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Albini, A. *Angew. Chem., Int. Ed.* **2011**, *50*, 1869. (b) Ravelli, D.; Albini, A.; Fagnoni, M. *Chem. Eur. J.* **2011**, *17*, 572. (c) Montanaro, S.; Ravelli, D.; Merli, D.; Fagnoni, M.; Albini, A. *Org. Lett.* **2012**, *14*, 4218. (d) Ryu, I.; Tani, A.; Fukuyama, T.; Ravelli, D.; Montanaro, S.; Fagnoni, M. *Org. Lett.* **2013**, *15*, 2554. (e) Orareya, H.; Ravelli, D.; Fagnoni, M.; Albini, A. *Adv. Synth. Catal.* **2013**, *355*, 2891. Also, see reviews: (f) Hill, C. L. *Synlett* **1995**, 127. (g) Tanielian, C. *Coord. Chem. Rev.* **1998**, *178-180*, 1165. (h) Hill, C. L. *J. Mol. Catal. A: Chem.* **2007**, *262*, 2. (i) Tzirakis, M. D.; Lykakis, I. N.; Orfanopoulos, M. *Chem. Soc. Rev.* **2009**, *38*, 2609.

(10) Okada, M.; Fukuyama, T.; Yamada, K.; Ryu, I.; Ravelli, D.; Fagnoni, M. *Chem. Sci.* **2014**, *5*, 2893.

(11) Roberts, B. P. *Chem. Soc. Rev.* **1999**, *28*, 25.

(12) See a review: Chatgililoglu, C.; Crich, D.; Komatsu, M.; Ryu, I. *Chem. Rev.* **1999**, *99*, 1991.

(13) (a) Texier, I.; Delaire, J. A.; Giannotti, C. *Phys. Chem. Chem. Phys.* **2000**, *2*, 1205. (b) Tanielian, C.; Coughon, F.; Seghrouchni, R. *J. Mol. Catal. A: Chem.* **2007**, *262*, 164.

(14) BDE of C-H bond for butane: primary C-H = 100.7 kcal/mol and secondary C-H = 98.3 kcal/mol; see: Luo, Y. -R. *Comprehensive Handbook of Chemical Bond Energies*; CRC Press: Boca Raton, FL, **2007**.

(15) Potter, A.; Tedder, J. *J. Chem. Soc., Perkin Trans. 2* **1982**, 1689.

(16) Recently,  $\gamma$ -selective fluorination of valeric acid methyl ester using TBADT was reported ( $\gamma/\beta = 81/19$ ). See: Halperin, S. D.; Fan, H.; Chang, S.; Martin, R. E.; Britton, R. *Angew. Chem., Int. Ed.* **2014**, *53*, 4690.

(17) For  $\gamma$ -preference to  $\beta$  in hydrogen abstraction of tetrahydropyran by *t*-butoxy radical. See, Jenkins, I. D. *J. Chem. Soc., Chem. Commun.* **1994**, 1227.